















- [36] N. Iblher, E. Gladilin, and B. G. Stark, "Soft-Tissue Mobility of the Lower Face Depending on Positional Changes and Age: A Three-Dimensional Morphometric Surface Analysis," *Plast Reconstr Surg*, vol. 131, no. 2, pp. 372–381, 2013.
- [37] E. Nkenke *et al.*, "Three-dimensional analysis of changes of the malar-midfacial region after LeFort I osteotomy and maxillary advancement," *Oral Maxillofac Surg*, vol. 12, no. 1, pp. 5–12, 2008.
- [38] J. Winzweig, A. Oliker, C. B. Cutting, S. J. Aston, and D. M. Smith, "Designing a Virtual Reality Model for Aesthetic Surgery," *Plast. Reconstr. Surg.*, vol. 116, no. 3, pp. 893–897, 2004.
- [39] C. Alfano, P. Mezzana, and N. Scuderi, "Acquisition and elaboration of superficial three-dimensional," *Indian J Plast Surg*, vol. 38, no. 1, pp. 22–27, 2005.
- [40] R. D. F. V. Dornelles, N. Alonso, L. A. L. Tissiani, A. R. Souza, and V. L. N. Cardim, "The use of a three-dimensional mesh in plastic surgery," *Rev. Bras. Cir. Plástica – Brazilian J. Plast. Surgery*, vol. 31, no. 1, pp. 25–31, 2016.
- [41] J. Y. Wong *et al.*, "Validity and Reliability of Craniofacial Anthropometric Measurement of 3D Digital Photogrammetric Images," *Cleft Palate-Craniofacial J.*, vol. 45, no. 3, pp. 232–239, 2007.
- [42] M. Van Der Vlis, K. M. Dentino, B. Vervloet, and B. L. Padwa, "Postoperative swelling after orthognathic surgery: A prospective volumetric analysis," *J. Oral Maxillofac. Surg.*, vol. 72, no. 11, pp. 2241–2247, 2014.
- [43] D. Y. S. Tanikawa, M. Aguená, D. F. Bueno, M. R. Passos-Bueno, and N. Alonso, "Fat grafts supplemented with adipose-derived stromal cells in the rehabilitation of patients with craniofacial microsomia," *Plast. Reconstr. Surg.*, vol. 132, no. 1, pp. 141–152, 2013.