

- Goto, R., Heinrich, K.M., Jitnarim, N., Suminski, R.R. et al. (2008). Obesity classification in military personnel: a comparison of body fat, waist circumference and body mass index measurements. *Mil Med.* 173, 67-73.
- Jebb, S.A., Cole, T.J., Doman, D., Murgatroyd, P.R., Prentice, A.M. (2000). Evaluation of the novel Tanitaq body-fat analyser to measure body composition by comparison with a four-compartment model. *Br J Nutr.* **83**: 115-122.
- Kissebah, A.H., Freedman, D.S., Peiris, A.N. (1989). Health risks of obesity. *Med Clin North Am.* 73 (1): 111–138.
- Kissebah, A.H., Vydellingum, N., Murray, R., et al. (1982). Relation of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab.* 4 (2): 254–260.
- Lemieux, S., Prud'homme, D., Tremblay, A., Bouchard, C., Despres, J.P. (1996). Anthropometric correlates to changes in visceral adipose tissue over 7 years in women. *Int J Obes Relat Metab Disord.* 20 (7): 618–624.
- Medina-Inojosa, J., Somers, V., Jenkins, S., Zundel, J., Johnson, L., et al. (2017). Validation of a White-light 3D Body Volume Scanner to Assess Body Composition. *Obes Open Access* 3(1):doi <http://dx.doi.org/10.16966/2380-5528.127>
- Mokdad, A.H., Ford, E.S., Bowman, B.A. et al. (2001). Prevalence of obesity, diabetes, and obesity-related health risk factors. *JAMA.* **289** (1), 76-79.
- National Family Health Survey (NFHS-4). 2015-16. National Fact Sheet. INDIA (Provisional data). International Institute for Population Sciences, Mumbai. Ministry of Health and Family Welfare. Government of India.
- Norafidah, A.R., Azmawati, M.N., and Norfazilah, A. (2013). Factors influencing abdominal obesity by waist circumference among normal BMI population. *Malaysian Journal of Public Health.* 13(1):37-47.
- Nguyen-Duy, T.B., Nichaman, M.Z., Church, T.S., Blair, S.N., Ross, R. (2003). Visceral fat and liver fat are independent predictors of metabolic risk factors in men. *Am J Physiol Endocrinol Metab.* 284 (6): E1065–E1071.
- Nunez, C., Gallagher, D., Visser, M., Pi-Sunyer, F.X., Wang, Z., Heymsfield, S.B. (1997). Bioimpedance analysis: evaluation of leg-to-leg system bases on pressure contact foot–pad electrodes. *Med Sci Sports Exerc.* **29**(4):524-531.
- Nuttall, Frank Q. (2015). Body Mass Index: Obesity, BMI, and Health A Critical Review. *Nutrition Today*: May/June 2015 - Volume 50 - Issue 3 - p 117–128.
- Pischon, T., Boeing, H., Hoffmann, K., et al. (2008). General and abdominal adiposity and risk of death in Europe. *N Engl J Med.* 359 (20): 2105–2120.
- Pouliot, M.C., Despres, J.P., Lemieux, S., et al. (1994). Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *Am J Cardiol.* 73 (7): 460–468.
- Pouliot, M.C., Despres, J.P., Nadeau, A., et al. (1990). Associations between regional body fat distribution, fasting plasma free fatty acid levels and glucose tolerance in premenopausal women. *Int J Obes.* 14 (4): 293–302.
- Prentice, A.M., Jebb, S.A. (2001). Beyond body mass index. *Obes Rev.* **2**(3):141-147.
- Singh, S.P., Sikri, G. and Garg, M.K. (2008). Body mass index and obesity: Tailoring “cut-off” for an Asian Indian male population. *MJAFI.* **64**, 350-353.
- Sundin, J., Fear, N.T., Wessely, S. and Rona, R.J. (2011). Obesity in the UK armed forces: Risk factors. *Military Medicine.* **176**(5), 507-512.
- Varte, L.R., Rawat, S., and Singh, I. (2013). Relationship of Body Mass Index, Waist Circumference and Waist-Stature Ratio with Body Fat of the Indian Gorkha Population. *Mal J Nutr* **19**(2): 185 – 192.
- Wajchenberg, B.L., Giannella-Neto, D., da Silva, M.E., Santos, R.F. (2002). Depot-specific hormonal characteristics of subcutaneous and visceral adipose tissue and their relation to the metabolic syndrome. *Horm Metab Res.* 34 (11–12): 616–621.
- Wang, J., Thornton, J.C., Russel, M., Burastero, S., Heymsfield, S. and Pierson, Jr .R.N. (1994). Asians have lower body mass index (BMI) but higher percent body fat than do whites: comparisons of anthropometric measurements. *The American Journal of Clinical Nutrition.* 60(1):23-28.
- Wildman, R.P., Muntner, P., Reynolds, K., et al. (2008). The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence and correlates of 2 phenotypes among the US population (NHANES 1999–2004). *Arch Intern Med.* 168 (15): 1617–1624.